

**Supplementary Table 2.** Comparison of donor factors according to median BMI value (BMI 22.7 kg/m<sup>2</sup>)

Factor	Low BMI (n = 342)	High BMI (n = 340)	p-value
Age, ≥65 yr	2 (0.6)	8 (2.4)	0.063
Male sex	176 (51.5)	198 (58.2)	0.076
BMI (kg/m <sup>2</sup> )			
<18.5 (underweight)	12 (3.5)	9 (2.6)	0.515
18.5–22.7 (normal)	130 (38.0)	105 (30.9)	0.050
22.7–25.0 (overweight)	93 (27.2)	108 (31.8)	0.190
25.0–30.0 (mildly obese)	90 (26.3)	100 (29.4)	0.367
≥30.0 (grossly obese)	17 (5.0)	18 (5.3)	0.848
eGFR (CKD-EPI) (mL/min/1.73 m <sup>2</sup> )	114.7 ± 12.5	113.5 ± 13.0	0.218
Kw/Rw ratio	3.4 ± 0.8	2.6 ± 0.7	<0.001

Data are expressed as number (%) or mean ± standard deviation.

BMI, body mass index; CKD-EPI, Chronic Kidney Disease-Epidemiology Collaboration; eGFR, estimated glomerular filtration rate; Kw/Rw, donated kidney weight/recipient body weight.