

Supplementary Table 3. Clinical characteristics at 48 hours after CRRT initiation according to Phoxilium use

Characteristic	All (n = 272)	Phoxilium (-) (n = 96)	Phoxilium (+) (n = 176)	p-value
SOFA_48hr	9.1 ± 3.9	8.0 ± 4.0	9.7 ± 3.7	0.001
White blood cell_48hr (10 ³ /μL)	13.5 ± 8.0	12.2 ± 6.8	14.2 ± 8.5	0.05
Hemoglobin_48hr (g/dL)	9.8 ± 2.1	10.1 ± 2.6	9.6 ± 1.7	0.05
RDW-CV_48hr (%)	15.5 ± 2.4	14.8 ± 2.0	15.8 ± 2.5	0.005
Hemoglobin_48hr/RDW-CV_48hr	0.6 ± 0.2	0.7 ± 0.2	0.6 ± 0.2	0.002
Albumin_48hr (g/dL)	2.8 ± 0.5	2.9 ± 0.6	2.8 ± 0.5	0.07
Potassium_48hr (mmol/L)	4.1 ± 0.7	4.1 ± 0.9	4.0 ± 0.6	0.91
Arterial pH_48hr	7.41 ± 0.09	7.40 ± 0.10	7.41 ± 0.08	0.93
Bicarbonate_48hr (mEq/L)	22.1 ± 4.6	23.1 ± 5.5	21.7 ± 4.0	0.03
BUN_48hr (mg/dL)	29.1 ± 17.2	34.5 ± 20.4	26.3 ± 14.4	<0.001
Creatinine_48hr (mg/dL)	1.8 ± 1.4	2.2 ± 2.0	1.5 ± 0.9	<0.001
Phosphorus_48hr (mg/dL)	3.3 ± 1.5	3.6 ± 2.0	3.1 ± 1.1	0.04
Calcium_48hr (mg/dL) ^a	8.2 ± 0.7	8.3 ± 0.8	8.2 ± 0.7	0.18

Data are expressed as mean ± standard deviation.

Phoxilium (-): the group of patients who did not receive Phoxilium within 48 hours after CRRT initiation. Phoxilium (+): the group of patients who received Phoxilium within 48 hours after CRRT initiation

BUN, blood urea nitrogen; CRRT, continuous renal replacement therapy; RDW-CV, red cell distribution width-coefficient of variation; SOFA, Sequential Organ Failure Assessment.

^aCorrected calcium (mg/dL) = measured total calcium (mg/dL) + 0.8×[4-measured serum albumin (g/dL)].