

Supplementary Table 4. The eGFR at dialysis initiation according to comorbidity index in symptomatic patients^a at dialysis initiation (n = 810)

Variable	Planned dialysis		Unplanned dialysis ^b		Total		p-value ^c
	No. (%)	eGFR (mL/min/1.73 m ²)	No. (%)	eGFR (mL/min/1.73 m ²)	No. (%)	eGFR (mL/min/1.73 m ²)	
Low-risk	175 (38.2)	4.93 ± 2.67	71 (20.2)	5.73 ± 3.71	246 (30.4)	5.16 ± 3.02	0.06
Medium-risk	250 (54.6)	6.85 ± 4.31	236 (67.1)	8.28 ± 5.30	486 (60.0)	7.54 ± 4.86	<0.001
High-risk	33 (7.2)	7.15 ± 2.91	45 (12.8)	10.06 ± 4.28	78 (9.6)	8.83 ± 4.00	<0.001

eGFR, estimated glomerular filtration rate.

^aSymptomatic patients: all patients except asymptomatic patients who started dialysis with progressive azotemia without specific symptoms. ^bUnplanned dialysis group was defined as starting maintenance dialysis in a life-threatening situation regardless of a permanent access device in place. ^cWe compared eGFR at dialysis initiation between the planned dialysis group and unplanned dialysis group.