

Supplementary Table 2. The percentage of cases of shared decision-making in clinical practice

Variable	No. of respondents	Mean \pm SE	p-value
Total	309	64.3 \pm 1.2	
Sex			0.94 ^a
Male	174	64.2 \pm 1.7	
Female	135	64.4 \pm 1.8	
Age (yr)			0.25 ^b
<30	90	63.4 \pm 2.1	
30–39	178	65.0 \pm 1.6	
40–49	26	61.0 \pm 4.7	
50–59	9	56.1 \pm 9.5	
60–65	6	80.0 \pm 7.3	
Position			0.22 ^b
Resident (1st/2nd grade)	148	63.4 \pm 1.6	
Resident (3rd/4th grade)	78	68.1 \pm 2.3	
Fellow	32	62.3 \pm 4.1	
Professor	51	62.0 \pm 3.5	
No. of patients treated by respondent (/wk)			0.75 ^b
<20	53	66.0 \pm 2.4	
20–49	144	64.1 \pm 1.7	
50–79	54	66.8 \pm 2.8	
80–99	18	55.8 \pm 7.9	
\geq 100	40	62.8 \pm 3.8	
No. of patients who made decisions ^c (/wk)			0.49 ^b
<2	207	63.5 \pm 1.5	
2–5	92	65.8 \pm 2.4	
5–7	5	60.0 \pm 4.5	
7–10	4	75.0 \pm 9.6	
\geq 10	1	60	

Data are expressed as number or mean \pm standard error.

The p-values obtained from ^aMann-Whitney test and ^bKruskal-Wallis test. ^cAccording to the Act on Decisions on Life-Sustaining Treatment for Patients in Hospice and Palliative Care or at the End-of-Life.