

**Supplementary Table 2. Inclusion and exclusion criteria**

KTR group		CKD group	
Inclusion criteria	Exclusion criteria	Inclusion criteria	Exclusion criteria
a. Adult patients (aged >18 years)	a. eGFR of <20 mL/min/1.73 m <sup>2</sup> or an eGFR decline >30% during the last 3 months	a. Adult patients (aged >18 years)	a. Chronic atrial fibrillation or other arrhythmias that could interfere with proper ABPM recording
b. Patients with kidney failure that received a kidney transplant at least 3 months ago	b. Chronic atrial fibrillation or other arrhythmias that could interfere with proper ABPM recording	b. Patients with stage 1–4 CKD	b. Modification of antihypertensive treatment during the previous 6 weeks
	c. Presence of non-functional arteriovenous fistulae that could interfere with proper ABPM recording		c. Myocardial infarction, angina pectoris, or stroke during the last 3 months
	d. Modification of antihypertensive treatment during the previous 6 weeks		d. History of malignancy or any other condition with poor prognosis
	e. Myocardial infarction, angina pectoris, or stroke during the last 3 months		e. Pregnancy
	h. History of malignancy or any other condition with poor prognosis		
	g. Pregnancy		

ABPM, ambulatory blood pressure monitoring; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; KTRs, kidney transplant recipients.