

Supplementary Table 3. Reclassification tables for the predicted 5-year risk categories of <5%, 5% to <10%, and ≥10% by the addition of albuminuria and eGFR levels to conventional models

Observed event	Reclassification				NRI
	To lower risk category	No change	To higher risk category	Sum	
Primary composite outcome in CKD population					0.13766
Event (-)	178,010 (15.91)	789,204 (70.52)	57,468 (5.14)	1,024,682 (91.56)	
Event (+)	19,951 (1.78)	55,104 (4.92)	19,350 (1.73)	94,405 (8.44)	
Sum	197,961 (17.69)	844,308 (75.45)	76,818 (6.86)	1,119,087 (100.00)	
Primary composite outcome in the general population					0.1335
Event (-)	34,075 (2.87)	1,088,691 (91.59)	25,305 (2.13)	1,148,071 (96.58)	
Event (+)	3,151 (0.27)	30,505 (2.57)	6,963 (0.59)	40,619 (3.42)	
Sum	37,226 (3.13)	1,119,196 (94.15)	32,268 (2.71)	1,188,690 (100.00)	
Secondary composite outcome in CKD population					0.13492
Event (-)	139,880 (12.50)	853,457 (76.26)	47,678 (4.26)	1,041,015 (93.02)	
Event (+)	15,941 (1.42)	44,724 (4.00)	17,407 (1.56)	78,072 (6.98)	
Sum	155,821 (13.92)	898,181 (80.26)	65,085 (5.82)	1,119,087 (100.00)	
Secondary composite outcome in the general population					0.064339
Event (-)	16,773 (1.41)	1,134,059 (95.40)	11,533 (0.97)	1,162,365 (97.79)	
Event (+)	1,808 (0.15)	21,585 (1.82)	2,932 (0.25)	26,325 (2.21)	
Sum	18,581 (1.56)	1,155,644 (97.22)	14,465 (1.22)	1,188,690 (100.00)	
Major cardiovascular events in CKD population					0.035484
Event (-)	35,007 (3.13)	996,797 (89.07)	31,088 (2.78)	1,062,892 (94.98)	
Event (+)	4,079 (0.36)	46,626 (4.17)	5,490 (0.49)	56,195 (5.02)	
Sum	39,086 (3.49)	1,043,423 (93.24)	36,578 (3.27)	1,119,087 (100.00)	
Major cardiovascular events in the general population					0.029738
Event (-)	7,066 (0.59)	1,150,151 (96.76)	7,875 (0.66)	1,165,092 (98.01)	
Event (+)	755 (0.06)	21,601 (1.82)	1,242 (0.10)	23,598 (1.99)	
Sum	7,821 (0.66)	1,171,752 (98.58)	9,117 (0.77)	1,188,690 (100.00)	

Data are expressed as number (%).

The primary composite outcome was major cardiovascular events or kidney disease progression. The secondary composite outcome was major cardiovascular events or kidney failure. Major cardiovascular events were critical atherosclerotic cardiovascular disease, hospitalized heart failure, and cardiovascular death.

CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; NRI, net reclassification improvement.