

Supplementary Table 1. Prognosis of inflammatory cytokines in non-SAKI patients

Variable	Survivors (n = 63)	Non-survivors (n = 21)	p-value
Creatinine ($\mu\text{mol/L}$)	58 (48–75)	60 (56–75)	0.74
WBC ($\times 10^9/\text{L}$)	7.4 (5.3–9.6)	10.5 (6.9–13.7)	0.04
CRP (mg/L)	25.8 (5.8–58.2)	68.7 (26.3–128.1)	0.004
PCT (ng/mL)	0.08 (0.04–0.32)	0.20 (0.09–1.13)	0.008
IL-2 (pg/mL)	0.45 (0.30–0.80)	0.62 (0.20–0.85)	0.66
IL-4 (pg/mL)	0.34 (0.12–0.73)	0.55 (0.36–1.26)	0.03
IL-6 (pg/mL)	11.2 (4.7–30.7)	21.8 (11.2–98.4)	0.01
IL-17A (pg/mL)	2.2 (0.7–4.0)	3.5 (1.3–5.7)	0.04
TNF- α (pg/mL)	0.43 (0.17–0.79)	0.73 (0.37–1.01)	0.045
IFN- γ (pg/mL)	1.27 (0.74–3.41)	1.17 (0.96–2.31)	0.92

Data are expressed as median (interquartile range).

CRP, C-reactive protein; IFN- γ , interferon gamma; IL-2, interleukin-2; IL-4, interleukin-4; IL-6, interleukin-6; IL-17A, interleukin-17A; PCT, procalcitonin; SAKI, sepsis-associated acute kidney injury; TNF- α , tumor necrosis factor alpha; WBC, white blood cell count.