

Supplementary Table 2. HR of adjusted variables in Cox regression model for anuria and mortalities

Variables	Anuria		All-cause mortality		CV mortality	
	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
Model 2						
Female (%)	0.71 (0.43–1.16)	0.17	1.32 (0.88–1.98)	0.18	1.13 (0.68–1.89)	0.64
BMI (kg/m ²)	1.17 (1.06–1.29)	0.001	1.06 (0.98–1.15)	0.15	1.04 (0.94–1.15)	0.42
Diabetes (%)	-	-	1.85 (1.24–2.75)	0.002	1.86 (1.10–3.17)	0.02
CVD (%)	-	-	0.74 (0.49–1.12)	0.15	0.97 (0.55–1.72)	0.92
Hemoglobin (g/L)	0.99 (0.98–1.01)	0.34	0.99 (0.98–1.00)	0.11	-	-
Albumin (g/L)	0.95 (0.89–1.00)	0.07	0.96 (0.91–1.00)	0.04	-	-
DBP (mmHg)	1.00 (0.97–1.02)	0.79	0.98 (0.96–1.00)	0.03	0.97 (0.95–1.00)	0.04
BUN (mmol/L)	-	-	0.97 (0.92–1.03)	0.38	-	-
nPCR (g/kg/day)	2.03 (0.70–5.89)	0.19	0.96 (0.27–3.47)	0.95	0.63 (0.21–1.86)	0.40
RKF (mL/min/1.73 m ²)	0.84 (0.73–0.96)	0.01	1.06 (0.96–1.17)	0.28	1.10 (0.98–1.23)	0.10
Model 3						
Female (%)	1.70 (1.06–2.72)	0.03	1.18 (0.82–1.71)	0.37	0.93 (0.57–1.52)	0.79
BMI (kg/m ²)	1.16 (1.05–1.28)	0.004	1.08 (1.00–1.17)	0.05	1.07 (0.97–1.18)	0.16
Diabetes (%)	-	-	1.86 (1.25–2.76)	0.002	1.91 (1.13–3.24)	0.02
CVD (%)	-	-	0.75 (0.50–1.14)	0.18	1.01 (0.57–1.80)	0.96
Hemoglobin (g/L)	0.99 (0.98–1.01)	0.24	0.99 (0.98–1.00)	0.12	-	-
Albumin (g/L)	0.94 (0.89–1.00)	0.06	0.95 (0.91–1.00)	0.03	-	-
DBP (mmHg)	1.00 (0.98–1.02)	0.96	0.98 (0.96–1.00)	0.049	0.98 (0.96–1.01)	0.13
BUN (mmol/L)	-	-	0.95 (0.90–1.01)	0.08	-	-
nPCR (g/kg/day)	2.02 (0.67–6.11)	0.21	1.69 (0.47–6.10)	0.42	1.13 (0.37–3.52)	0.83
Urine (/100 mL increase)	0.96 (0.91–1.01)	0.14	0.98 (0.94–1.03)	0.45	0.95 (0.89–1.01)	0.09

BMI, body mass index; BUN, blood urea nitrogen; CI, confidence interval; CV, cardiovascular; CVD, cardiovascular disease; DBP, diastolic blood pressure; HR, hazard ratio; nPCR, normalized protein catabolic rate; RKF, residual kidney function.