

Supplementary Table 3. Univariate and multivariate analyses of risk factors for death-censored graft failure

Variable	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Recipient age	1.04 (1.01–1.07)	0.02	1.03 (1.00–1.06)	0.07
Male recipient	2.51 (1.15–5.48)	0.02	2.28 (1.04–5.01)	0.04
Recipient BMI	1.03 (0.94–1.13)	0.49		
HLA class I mismatch \geq 3	1.71 (0.91–3.21)	0.09		
HLA class II mismatch \geq 2	1.52 (0.76–3.61)	0.24		
Male donor	1.41 (0.72–2.74)	0.31		
Donor BMI	1.02 (0.93–1.12)	0.72		
Donor type				
Living	1 (Reference)		1 (Reference)	
SCD	1.85 (0.86–3.98)	0.12	2.07 (0.95–4.49)	0.07
ECD	2.56 (1.18–5.53)	0.02	2.07 (0.93–4.62)	0.08
CMV antigenemia	0.86 (0.45–1.66)	0.65		
BK viremia	1.77 (0.32–1.84)	0.56		
Induction immunosuppression		0.66		
Basiliximab	1 (Reference)			
r-ATG	1.16 (0.60–2.26)			
Rejection episode	2.39 (1.24–4.60)	0.009	2.56 (1.32–4.96)	0.006

BMI, body mass index; CI, confidence interval; CMV, cytomegalovirus; ECD, extended criteria donor; HLA, human leukocyte antigen; HR, hazard ratio; r-ATG, rabbit anti-thymocyte globulin; SCD, standard criteria donor.