

Supplementary Table 5. Univariate and multivariate analyses of risk factors for patient death due to infection

Variable	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Recipient age	1.18 (1.09–1.27)	<0.001	1.14 (1.05–1.24)	0.002
Male recipient	2.60 (0.56–12.05)	0.22		
Recipient BMI	1.15 (0.99–1.34)	0.07		
Donor age	1.09 (1.03–1.14)	0.24	1.00 (0.94–1.06)	0.99
Male donor	0.41 (0.12–1.40)	0.16		
Donor BMI	1.15 (0.98–1.35)	0.09		
Donor type				
Living	1 (Reference)		1 (Reference)	
SCD	3.76 (0.34–41.45)	0.28	3.13 (0.27–35.86)	0.36
ECD	18.72 (2.34–149.83)	0.006	5.97 (0.48–74.66)	0.17
Induction immunosuppression		0.66		0.50
Basiliximab	1 (Reference)		1 (Reference)	
r-ATG	1.16 (0.60–2.26)		1.79 (0.33–9.79)	
Rejection episode	0.50 (0.13–1.89)	0.31		

CI, confidence interval; BMI, body mass index; ECD, extended criteria donor; HR, hazard ratio; r-ATG, rabbit anti-thymocyte globulin; SCD, standard criteria donor.