

Supplementary Table 6. HRs for all-cause mortality, MACEs, and composite outcome according to the Framingham risk score (patient with living-related donor kidney transplant, n = 3,135)

Outcome	Univariable analysis		Multivariable analysis	
	HR (95% CI)	p-value	HR (95% CI)	p-value
All-cause mortality		0.02		0.17
Q1	Reference		Reference	
Q2	0.84 (0.19–3.74)	0.82	0.30 (0.03–2.93)	0.30
Q3	3.80 (1.24–11.66)	0.02	2.51 (0.64–9.85)	0.19
Q4	2.98 (0.92–9.69)	0.07	2.08 (0.46–9.38)	0.34
MACE		<0.001		0.016
Q1	Reference		Reference	
Q2	10.15 (1.29–80.09)	0.03	5.55 (0.64–48.11)	0.12
Q3	20.38 (2.71–153.11)	0.03	8.40 (1.00–70.28)	0.05
Q4	37.27 (5.06–274.34)	<0.001	19.36 (2.32–161.62)	0.006
Composite outcome		<0.001		0.13
Q1	Reference		Reference	
Q2	1.18 (0.64–2.18)	0.60	0.78 (0.35–1.74)	0.55
Q3	1.85 (1.06–3.25)	0.03	1.26 (0.60–2.65)	0.54
Q4	2.99 (1.76–5.09)	<0.001	1.94 (0.89–4.20)	0.095

Multivariable analysis includes covariates of donor type, primary causes of kidney disease, body mass index, desensitization, serum phosphorus levels, and donor-specific antigens.

CI, confidence interval; HR, hazard ratio; MACE, major adverse cardiovascular events; Q, Framingham risk score quartile.