

Supplementary Table 7. HRs for all-cause mortality, MACEs, and composite outcome according to the Framingham risk score (patient with deceased donor kidney transplant, n = 1,547)

Outcome	Univariable analysis		Multivariable analysis	
	HR (95% CI)	p-value	HR (95% CI)	p-value
All-cause mortality		<0.001		0.01
Q1	Reference		Reference	
Q2	0.79 (0.23–2.72)	0.71	0.81 (0.20–3.26)	0.77
Q3	2.57 (0.95–6.96)	0.06	2.04 (0.62–6.65)	0.24
Q4	5.57 (2.20–14.11)	<0.001	4.05 (1.28–12.79)	0.02
MACE	MACE	0.001		0.22
Q1	Reference		Reference	
Q2	1.78 (0.55–5.79)	0.34	2.03 (0.41–10.15)	0.39
Q3	3.66 (1.25–10.77)	0.02	3.13 (0.66–14.89)	0.15
Q4	5.50 (1.94–15.62)	0.001	4.36 (0.92–20.66)	0.06
Composite outcome		<0.001		0.03
Q1	Reference		Reference	
Q2	0.95 (0.53–1.72)	0.87	0.96 (0.41–2.21)	0.92
Q3	1.31 (0.76–2.28)	0.33	1.02 (0.44–2.37)	0.97
Q4	2.39 (1.45–3.94)	0.001	2.29 (1.03–5.06)	0.04

Multivariable analysis includes covariates of donor type, primary causes of kidney disease, body mass index, desensitization, serum phosphorus levels, and donor-specific antigens.

CI, confidence interval; HR, hazard ratio; MACE, major adverse cardiovascular events; Q, Framingham risk score quartile.