

**Supplementary Table 2.** Incidence rate and HRs of decreased renal function and proteinuria in donors with hypertension and normotensive donors after transplantation in the entire cohort

Variable	No. of events (%)	Person-year	Incidence rate <sup>a</sup>	Unadjusted HR (95% CI)	p-value	Adjusted <sup>b</sup> HR (95% CI)	p-value
Mild decrease in renal function							
Normotensive donors	491 (10.1)	82,980	5.9	Reference		Reference	
Hypertensive donors	104 (16.2)	10,200	10.2	1.73 (1.40–2.14)	<0.001	0.87 (0.70–1.09)	0.22
Moderate decrease in renal function							
Normotensive donors	33 (0.7)	82,356	0.4	Reference		Reference	
Hypertensive donors	15 (2.3)	9,900	1.5	3.75 (2.04–6.91)	<0.001	1.52 (0.79–2.94)	0.21
Proteinuria							
Normotensive donors	113 (2.3)	79,584	1.4	Reference		Reference	
Hypertensive donors	23 (3.6)	9,780	2.4	1.69 (1.08–2.64)	0.02	1.77 (1.10–2.85)	0.02

CI, confidence interval; HR, hazard ratio.

<sup>a</sup>Events per 1,000 person-year. <sup>b</sup>All analyses are adjusted for the following covariates: age, sex, body mass index, diabetes mellitus, previous cardiovascular disease, and predonation estimated glomerular filtration rate.