

Supplementary Table 9. HRs for 28- and 90-day mortality based on PAR quartiles

Mortality	Model 1 ^a		Model 2 ^b		Model 3 ^c	
	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
28-Day						
Q1	Reference		Reference		Reference	
Q2	0.75 (0.64–0.89)	<0.001	0.74 (0.63–0.88)	<0.001	0.74 (0.63–0.88)	<0.001
Q3	0.59 (0.50–0.71)	<0.001	0.58 (0.48–0.69)	<0.001	0.58 (0.48–0.70)	<0.001
Q4	0.54 (0.45–0.64)	<0.001	0.52 (0.43–0.63)	<0.001	0.53 (0.44–0.63)	<0.001
90-Day						
Q1	Reference		Reference		Reference	
Q2	0.78 (0.66–0.91)	0.002	0.77 (0.65–0.90)	0.002	0.77 (0.65–0.91)	0.002
Q3	0.61 (0.51–0.72)	<0.001	0.59 (0.50–0.71)	<0.001	0.60 (0.50–0.71)	<0.001
Q4	0.55 (0.46–0.66)	<0.001	0.54 (0.45–0.64)	<0.001	0.54 (0.46–0.65)	<0.001

CI, confidence interval; HR, hazard ratio; PAR, platelet-to-albumin ratio.

^aUnadjusted model; ^badjusted for age and sex; ^cmodel with additional adjustments for Charlson Comorbidity Index score, and body mass index.